arXiv:cond-mat/9907137v1 [cond-mat.str-el] 9 Jul 1999

Hidden non-Fermi liquid behavior due to crystal field quartet

Mikito Koga
Department of Physics, University of California Davis, CA 95616
(February 1, 2008)

We study a realistic Kondo model for crystal field quartet ground states having magnetic and non-magnetic (quadrupolar) exchange couplings with conduction electrons, using the numerical renormalization group method. We focus on a local effect dependent on singlet excited states coupled to the quartet, which reduces the non-magnetic coupling significantly and drives non-Fermi liquid behavior observed in the calculated quadrupolar susceptibility. A crossover from the non-Fermi liquid state to the Fermi liquid state is characterized by a small energy scale very sensitive to the non-magnetic coupling. On the other hand, the Kondo temperature observed in the magnetic susceptibility is less sensitive. The different crystal-field dependence of the two exchange couplings may be related to the different x dependence of quadrupolar and magnetic ordering temperatures in $Ce_xLa_{1-x}B_6$.

PACS numbers: 75.20.Hr, 71.10.Hf, 71.27.+a, 72.15.Qm

I. INTRODUCTION

The importance of orbital degrees of freedom has been emphasized for heavy fermion systems such as Ce or U based compounds.[1–3] Since the f-electrons have localized character, the crystal-field multiplets play important roles in the low temperature physics. The multichannel Kondo effect due to the f-shell impurities is one of the possible origins of the non-Fermi liquid (NFL) behavior in the heavy fermion metals.[1] The expected logarithmic divergence in their magnetic susceptibility χ and specific heat coefficient γ with decreasing temperatures is observed in the dilute limit of the f-shell ions in $U_xTh_{1-x}Ru_2Si_2$[4] or $Ce_xLa_{1-x}Pd_2Al_3$.[5] This can be explained by the theory of the two-channel Kondo (2CK) effect for each case.[6] For the U case, the NFL behavior is due to a non-Kramers doublet in f^2, while for the Ce case, it is associated with a Kramers doublet in f^1 coupled to a non-Kramers doublet in f^2. These non-Kramers doublets give rise to two independent and equivalent exchange interactions between the conduction electrons and the f-shell moments. The exchange coupling is quadrupolar for U, while it is magnetic (dipolar or octupolar) for Ce. We can also apply a similar scenario to account for a power-low temperature dependent χ and γ in $UCu_{5-x}Pd_x$.[7,8] This NFL behavior may be caused by a quadrupolar Kondo coupling due to a triplet ground state of U.[9]

Our next interest in this work is what type of Kondo effect can occur for a quartet f-electron state in realistic metals. The quartet is coupled to the conduction electrons via a quadrupolar exchange coupling as well as a dipolar one. An octupolar exchange interaction has to be taken into account, too, since it has comparable amplitude to the others. The Kondo effect originating from the quartet is realized in $Ce_xLa_{1-x}B_6$, where the ground Γ_8 quartet is separated very well from the excited Γ_7 doublet by 540K, implying that the admixture of these states can be neglected at low temperatures. The observed χ, γ, resistivity, and thermopower in this alloy have been described by the Coqblin-Schrieffer (CS) Model with $SU(4)$ symmetry.[10] This model is derived from an Anderson model for orbitally degenerate conduction electrons and impurities where the valence fluctuation is restricted to f^0-f^1 and the Coulomb interaction on the impurity site is taken to be infinity. In fact, the $SU(4)$ symmetry is broken in real systems because of the f^2 configuration. To confirm the robustness of the CS model, we have to take into account more generalized exchange interactions associated with the atomic structure.[11] Pavarini and Andreani studied the stability of $SU(N)$ symmetry against some anisotropic exchange interactions based on their spherically symmetric model.[12] They concluded that the local Fermi-liquid (FL) fixed point derived from the CS model is always stable and the Kondo temperature is reduced. The recent study of Kusunose and Kuramoto on the same type of Kondo model found that an NFL state is realized in the absence of non-magnetic exchange couplings and is destabilized by the particle-hole symmetry breaking.[13]

We cannot find any evidence of NFL behavior in the existing experimental data for $Ce_xLa_{1-x}B_6$. It is not easy to obtain direct information of Ce single-site effects for dense Ce cases since the long-range order occurs below 3.3K, very close to the Kondo temperature ($\sim$ 2K). However, we note the difference between magnetic and quadrupolar effects found in the phase diagram dependent on the Ce concentration x. The transition temperature of the antiferro-quadrupolar order T_{Q} depends on x more sensitively than the Néel temperature T_{N}.[14,15] The former decreases very rapidly with decreasing x. We can propose two possible scenarios to explain this behavior: (1) the quadrupolar order is destabilized more easily by the lattice disorder intensified with decreasing x, and (2) the quadrupolar exchange coupling between the conduction electrons and the Γ_8 quartet becomes smaller than the magnetic one, which results in the rapid decrease of the quadrupolar RKKY coupling. Since the latter has not been considered before, we stress that the strong suppression of the quadrupolar exchange coupling is due to the crystal-field states of Ce varying with decreasing x from $x = 1$. It is also interesting that this quadrupolar property is very important for the possibility of a new NFL state as discussed later.

In this paper we present a realistic Kondo model with a spherical tensor form. By using a Schrieffer-Wolff transformation,[16] it is derived from an Anderson model with an orbitally degenerate impurity on which the f^1 Γ_8 quartet is coupled to the f^0 configuration and the f^2 Γ_1 singlet via the hybridization with the conduction electrons. The model consists of a magnetic exchange term (dipolar and octupolar interactions are combined) and a non-magnetic term (a potential scattering and a quadrupolar exchange interaction are combined). If we tune up the f-electron energy levels to leave the magnetic part alone, we can obtain an NFL state corresponding exactly to that of the 2CK model, except for the conversion of the charge and spin indices. First, we discuss this NFL state realized only in a particle-hole symmetric case, based on the numerical renormalization group (NRG) analysis.[17] Next, we investigate the realistic case where the particle-hole symmetry is broken and the effect of our NFL fixed point is observed in the quadrupolar (strain) susceptibility. The most important result is the existence of two characteristic energy scales: one corresponds to the Kondo temperature associated with the CS limit, while the other characterizes a crossover to the FL state at lower temperatures. The latter new scale is connected to the NFL fixed point and can be very small when the quadrupolar exchange coupling is small. This quadrupolar coupling destabilizes the NFL state in the same manner as a spin field in the 2CK model.[18]

The remainder of this paper is organized as follows. In Sec. II, we explain the different crystal-field dependence of the magnetic and non-magnetic exchange couplings and describe how to calculate magnetic and quadrupolar susceptibilities for the impurity with the NRG method. In Sec. III, we discuss the FL and NFL states based on the NRG results. Finally, Sec. IV gives the concluding remarks and discussions with some proposals for experiments.

II. MODEL

As mentioned above, we restrict the f-electron states to f^0, an f^1 Γ_8 quartet, and an f^2 Γ_1 singlet in our Anderson model. These are possible ground states, accessed by varying two parameters of a crystal-field Hamiltonian.[19] Here we assume that the energies of the excited states in each configuration are so large as to be neglected at low temperatures. Only in this case can we map a derived Kondo model to an exchange model expressed with a spherical tensor form. Applying a Schrieffer-Wolff transformation to the f^0-f^1 valence fluctuation, we obtain the CS model with $SU(4)$ symmetry. To rewrite this Hamiltonian with spherical tensor operators, we express the fourfold Γ_8 states in terms of J_z of a $J = 3/2$ spin operator in the following way:[20]

$$|\Gamma_{8,3/2}\rangle = -\sqrt{\frac{1}{6}}|+\frac{3}{2}\rangle - \sqrt{\frac{5}{6}}|-\frac{5}{2}\rangle, \tag{1}$$

$$|\Gamma_{8,1/2}\rangle = |+\frac{1}{2}\rangle, \tag{2}$$

$$|\Gamma_{8,-1/2}\rangle = -|-\frac{1}{2}\rangle, \tag{3}$$

$$|\Gamma_{8,-3/2}\rangle = \sqrt{\frac{1}{6}}|-\frac{3}{2}\rangle + \sqrt{\frac{5}{6}}|+\frac{5}{2}\rangle, \tag{4}$$

where we assume that these states are constructed from the single-electron states with the total angular momentum $j = 5/2$ because of a strong spin-orbit coupling in Ce, and $|M\rangle$ ($M = -5/2, -3/2, \cdots, 5/2$) represents an eigenstate of j_z. The CS Hamiltonian can be expanded with respect to scalar products of two kinds of irreducible tensor operators with rank p, and the corresponding effective exchange Hamiltonian is given by

$$H_{\mathrm{ex},10} = I_{10} \sum_{kk'mm'} a^\dagger_{k'm'} a_{km} \times \left[\frac{1}{5}(T^{(1)}_{m'm} + \frac{4}{9}T^{(3)}_{m'm}) + \frac{1}{4}(1 + \frac{4}{9}T^{(2)}_{m'm}) \right], \tag{5}$$

where

$$T^{(p)}_{m'm} = \sum_{q=-p}^{p} (-1)^q \left(j^{(p)}_{-q}\right)_{m'm} J^{(p)}_q \tag{6}$$

represents a dipolar, a quadrupolar, and an octupolar exchange interaction for $p = 1$, $p = 2$, and $p = 3$, respectively. Here $a^\dagger_{km}$ (a_{km}) is a creation (annihilation) operator for a conduction electron with wave vector k and orbital m. The tensor $j^{(p)}_q$ ($J^{(p)}_q$) is constructed from a $j = 3/2$ ($J = 3/2$) spin operator for the fourfold Γ_8 conduction electrons (the Γ_8 quartet). For each p, the tensor operator has $(2p+1)$ components and is derived from the following equations,

$$J^{(p)}_p = (-1)^p \sqrt{\frac{(2p-1)(2p-3)\cdots 3\cdot 1}{2p(2p-1)\cdots 2}} J^p_+, \tag{7}$$

$$J^{(p)}_{q-1} = \frac{1}{\sqrt{(p+q)(p-q+1)}}[J_-, J^{(p)}_q]. \tag{8}$$

We take the Fermi energy as the origin of energies, and the coupling constant I_{10} is given by $V_{10}^2/\Delta E_{10}$, where V_{10} is the mixing parameter for f^0-f^1 and ΔE_{10} is the energy level of f^0 measured from that of the f^1 Γ_8 quartet. The valence fluctuation from f^1-f^2 also leads to the similar form of the effective exchange Hamiltonian since the twofold Γ_7 ($j = 5/2$) conduction electrons cannot contribute to this exchange process, being restricted by the Γ_1 symmetry ($\Gamma_1 = \Gamma_6 \otimes \Gamma_6 \oplus \Gamma_7 \otimes \Gamma_7 \oplus \Gamma_8 \otimes \Gamma_8$) of the intermediate state in f^2. The SU(4) symmetry is also conserved in the exchange interaction, and we obtain the corresponding Hamiltonian for f^1-f^2,

$$H_{\mathrm{ex},12} = I_{12} \sum_{kk'mm'} a^\dagger_{k'm'} a_{km} \times \left[\frac{1}{5}(T^{(1)}_{m'm} + \frac{4}{9} T^{(3)}_{m'm}) - \frac{1}{4}(1 + \frac{4}{9} T^{(2)}_{m'm}) \right], \quad (9)$$

where the sign of the potential scattering and quadrupolar coupling terms is negative, while it is positive in the Hamiltonian (5). Here the coupling constant I_{12} is given by $V_{12}^2/\Delta E_{12}$, where the definition of each parameter follows that for I_{10}. If the $\Gamma_8 \otimes \Gamma_8$ part of the f^2 Γ_1 state consists of single-electron states given in (1)–(4), we can obtain $V_{12}/V_{10} = 1/\sqrt{6}$ which is related to the Clebsch-Gordan coefficients. After we put the two Hamiltonians (5) and (9) together, we obtain the complete exchange Hamiltonian with two exchange couplings as follows:

$$H_{\mathrm{ex}} = \sum_{kk'mm'} a^\dagger_{k'm'} a_{km} (I_{\mathrm{o}} T^{(\mathrm{o})}_{m'm} + I_{\mathrm{e}} T^{(\mathrm{e})}_{m'm}), \quad (10)$$

where

$$T^{(\mathrm{o})}_{m'm} = \frac{1}{5}(T^{(1)}_{m'm} + \frac{4}{9} T^{(3)}_{m'm}), \quad (11)$$

$$T^{(\mathrm{e})}_{m'm} = \frac{1}{4}(1 + \frac{4}{9} T^{(2)}_{m'm}), \quad (12)$$

$$\frac{I_{\mathrm{e}}}{I_{\mathrm{o}}} = \frac{1 - (I_{12}/I_{10})}{1 + (I_{12}/I_{10})} \quad (\equiv \alpha). \quad (13)$$

Here I_{o} is always positive and antiferromagnetic, while the sign of I_{e} can change. We note that the transformation $I_{\mathrm{e}} \leftrightarrow -I_{\mathrm{e}}$ only causes the conversion between particles and holes. The ratio of the couplings α represents the anisotropy apart from the $SU(4)$ symmetry ($\alpha = 1$). As shown in Eq. (13), the quadrupolar coupling can be very small depending on the exchange couplings I_{10} and I_{12}, i.e., on the valence fluctuations f^0-f^1 and f^1-f^2. Thus the quadrupolar coupling is suppressed by the local environment more strongly than the dipolar and octupolar couplings. Recently Kusunose and Kuramoto studied the same type of Kondo model for arbitrary ranks of tensors by the perturbation renormalization group analysis.[13] They found that an unstable NFL fixed point is realized when all the exchange interactions expressed by $j^{(2n)}_{-q} J^{(2n)}_q$ (n is integer) vanish and particle-hole symmetry is conserved. This corresponds to the case $I_{\mathrm{e}} = 0$ ($I_{10} = I_{12}$) in our realistic Kondo model. This is realized when $\Delta E_{12}/\Delta E_{10}$ is exactly equal to $(V_{12}/V_{10})^2$ ($= 1/6$) for the crystal-field energies of f-electrons. Such complete cancellation of I_{10} and I_{12} is unlikely in real systems, but the effect of the NFL state should be observed at finite temperatures as discussed later.

To investigate our Kondo model given by

$$H = \sum_{km} \varepsilon_k a^\dagger_{km} a_{km} + H_{\mathrm{ex}}, \quad (14)$$

where ε_k is the kinetic energy of the conduction electron, we transform the Hamiltonian to a hopping Hamiltonian for the NRG calculation. Following Wilson,[17] we obtain the recursion relation

$$H_{N+1} = \Lambda^{1/2} H_N + (f^\dagger_{N+1,m} f_{Nm} + \mathrm{H.c.}), \quad (15)$$

and

$$H_0 = \Lambda^{-1/2} \sum_{mm'} f^\dagger_{0,m'} f_{0,m} (\tilde{I}_{\mathrm{o}} T^{(\mathrm{o})}_{m'm} + \tilde{I}_{\mathrm{e}} T^{(\mathrm{e})}_{m'm}), \quad (16)$$

for the impurity, where through the Fourier transformation $f^\dagger_{nm}$ (f_{nm}) is derived from $a^\dagger_{km}$ (a_{km}) defined in the logarithmically descretized conduction band, and $\tilde{I}_{\mathrm{o(e)}}$ is equal to $2I_{\mathrm{o(e)}}\rho/(1+\Lambda^{-1})$, where ρ is the density of states of the conduction electrons at the Fermi sea. In order to obtain the temperature dependent magnetic (χ) and quadrupolar (χ_{Q}) susceptibilities for the local Γ_8 quartet, we calculate the magnetization M_z and quadrupolar response Q_{zz}, respectively, to very small local external fields. For the latter, the lattice is distorted along a c-axis. In the NRG calculation, M_z and Q_{zz} are given by

$$M_z = \frac{\mathrm{Tr} J_z \exp(-\bar{\beta} H'_N)}{\mathrm{Tr} \exp(-\bar{\beta} H'_N)}, \quad (17)$$

$$Q_{zz} = \frac{\mathrm{Tr}(\lambda O^2_0) \exp(-\bar{\beta} H''_N)}{\mathrm{Tr} \exp(-\bar{\beta} H''_N)}. \quad (18)$$

Here O^2_0 is defined by $[3J^2 - J(J+1)]$ and $\bar{\beta}$ (~ 1) is related to the physical temperature

$$k_{\mathrm{B}} T/D = \frac{1 + \Lambda^{-1}}{2} \Lambda^{-(N-1)/2} / \bar{\beta}, \quad (19)$$

where D is a half width of the conduction band. For the Hamiltonian H'_N in Eq. (17), the Zeeman term $-g_J \mu_{\mathrm{B}} J_z h_z$ ($g_J = 6/7$ is a Landé's g factor for the Γ_8 quartet and h_z is a local magnetic field) is added to the exchange Hamiltonian H_0. For H''_N in Eq. (18), the similar term $-\lambda O^2_0 h_q$ (h_q is a local quadrupolar field) is added to H_0, where we determine λ to satisfy

$$\langle \Gamma_{8,\pm 3/2} | \lambda O^2_0 | \Gamma_{8,\pm 3/2} \rangle = +1, \quad (20)$$

$$\langle \Gamma_{8,\pm 1/2} | \lambda O^2_0 | \Gamma_{8,\pm 1/2} \rangle = -1. \quad (21)$$

The susceptibilities χ and χ_{Q} are given by M_z/h_z and Q_{zz}/h_q, respectively, if the local external fields are small enough. Throughout this paper we take k_{B}, μ_{B} and D to be unity. In the NRG calculation, we take $\Lambda = 3$ and keep about 800 lowest-lying states at each renormalization step.

III. RESULTS

Let us begin with the case $\alpha = 0$ in (13). Because of the particle-hole symmetry, we can express eigenstates with an axial charge operator $\vec{q}$ defined by[21]

$$q_+ = \sum_{n=0}^{\infty}(f^\dagger_{n,3/2}f^\dagger_{n,-3/2} - f^\dagger_{n,1/2}f^\dagger_{n,-1/2}), \qquad (22)$$

$$q_z = \frac{1}{2}\sum_{n=0}^{\infty}\sum_{m}(f^\dagger_{nm}f_{nm} - \frac{1}{2}). \qquad (23)$$

This operator satisfies the $SU(2)$ Lie algebra,

$$[q_z, q_\pm] = \pm q_\pm, \quad [q_+, q_-] = 2q_z. \qquad (24)$$

After a large number of renormalization steps N, we reach an NFL fixed point where $H_N = H_{N+2}$ is satisfied. Each energy level is labeled by (q, j) where j is the total angular momentum. As shown in Table I (a), the ground state has twofold degeneracy related to the particle-hole symmetry ($q_z = \pm 1/2$) and no angular momentum ($j = 0$). The lowest energy spectrum is same as that of NFL fixed point for the 2CK model. The eigenstates for the latter are given in (b). The energies in Table I are derived by the boundary conformal field theory (CFT)[22] which is applied to the 2CK model with the $U(1) \times SU(2) \times SU(2)$ symmetry. Here $v_{\rm F}$ is the Fermi velocity and l is the system size. Each energy is equal to the corresponding NRG energy multiplied by a factor ($\simeq 5/8$) for $\Lambda = 3$ case.[18] Our Kondo model with particle-hole symmetry has the $SU(2)$ symmetry for both q and j, and each of the four lowest-lying eigenstates in (a) satisfies the following equation

$$E = \frac{v_{\rm F}\pi}{l}\left[\frac{q(q+1)}{4} + \frac{j(j+1)}{12}\right], \qquad (25)$$

for the corresponding energy. Each energy in the last column is measured from that of the ground state ($= 3/16$). These eigenstates can be connected to the primary states in the CFT. The CFT gives the same equation to NFL fixed-point energies for a Kondo model with only dipolar exchange interaction between $j = 3/2$ conduction electrons and a $J = 1/2$ local spin.[21] This indicates applicability of the CFT to our Kondo model with the octupolar exchange interaction, although it is impossible to absorb the impurity spin in the conduction electron current.

The new NFL fixed point is stabilized by the octupolar exchange interaction $T^{(3)}$ in the Hamiltonian (10). On the other hand, the dipolar exchange interaction $T^{(1)}$ leads to another NFL fixed point if both $T^{(2)}$ and $T^{(3)}$ are neglected. The lowest NRG energy levels in this case are same as those for the Kondo model with exchange interaction between $j = 3/2$ conduction electrons and a $J = 1/2$ local spin, if even and odd numbers of the NRG iteration step N are converted each other. The NFL fixed point due to $T^{(1)}$ is stable against the potential scattering and is destabilized by $T^{(2)}$ and $T^{(3)}$. If only $T^{(1)}$ and $T^{(3)}$ are left in the exchange interaction and these couplings are varied independently, we can obtain the competition between each other. For the exchange Hamiltonian (10), we always reach the NFL fixed point stabilized by $T^{(3)}$ when we fix $\alpha = 0$.

The particle-hole symmetry is usually broken ($\alpha \neq 0$) and the quadrupolar coupling exists in any real systems. In Table I, the axial charge q for the new NFL fixed point (a) completely corresponds to the spin S for the 2CK fixed point (b). As a spin field in the 2CK model, the potential scattering and quadrupolar exchange terms destabilize the new NFL fixed point, and leads to the $SU(4)$ symmetric FL fixed point realized for the CS model. The crossover from the NFL state to the FL state is related to the recovery of the $SU(4)$ symmetry. In the NRG calculation, the doublet ground state ($q = 1/2, j = 0$) for $\alpha = 0$ is lifted by the coupling $I_{\rm e}$. The NRG energy flow diagram shows that one of the separated states becomes unstable as the renormalization step N increases, and finally merges to a fivefold degenerate excited state with $j = 2$. We define the crossover temperature $T^*_{\rm s}$ as the number of the NRG step at which the energy difference between the two states reach less than 0.01. Figure 1 shows that $T^*_{\rm s}$ is proportional to α^2, similar to a crossover temperature vs. a spin field for the 2CK model.[18]

Next we discuss the crossover to the FL state by calculating $\chi(T)$ and $\chi_{\rm Q}(T)$. In the CS limit ($\alpha = 1$), these susceptibilities show the same temperature dependence. If they are normalized by the $T = 0$ values, we can obtain the complete agreement with each other. The difference is enhanced as the symmetry of the exchange interaction is lowered from the $SU(4)$ symmetry, i.e, α decreases from unity. In Fig. 2, $\chi_{\rm Q}(T)$ for $\alpha = 0$ behaves logarithmically down to $T = 0$ as does the spin susceptibility for the 2CK model. Based on this logarithmic curve, we can estimate the Kondo temperature $T^{(0)}_{\rm K}$ as a function of the magnetic coupling $I_{\rm o}$. Once the non-magnetic coupling $I_{\rm e}$ is introduced ($\alpha \neq 0$), the FL state appears below $T^*_{\rm s}$ and $\chi_{\rm Q}$ becomes constant. The $T = 0$ value of $\chi_{\rm Q}$ decreases as α increases. As shown in Fig. 3, $\chi_{\rm Q}(T = 0) \propto \ln\alpha$ is satisfied for the small values of α. In analogy with the spin susceptibility for the two-channel Kondo model with channel anisotropy,[23,24] $\chi_{\rm Q}$ is scaled by $T^{(0)}_{\rm K}$ and $T_{\rm s}$: the latter is proportional to $T^*_{\rm s}$. Since $T_{\rm s}$ is proportional to α^2, we can express $\chi_{\rm Q}$ with the following equation

$$\chi_{\rm Q} = \frac{a}{T^{(0)}_{\rm K}}\ln\left[\frac{T^{(0)}_{\rm K}}{\max(T_{\rm s}, T)}\right], \qquad (26)$$

for $T_{\rm s} \ll T^{(0)}_{\rm K}$. We obtain $(a, T^{(0)}_{\rm K}, T_{\rm s}/\alpha^2) = (0.60, 2.6 \times 10^{-3}, 8.1 \times 10^{-3})$ for $\tilde{I}_{\rm o} = 0.2$ and $(0.62, 3.1 \times 10^{-2}, 6.1 \times 10^{-2})$ for $\tilde{I}_{\rm o} = 0.3$.

These two energy scales are also found in χ, which we conclude from the detailed analysis shown below. For the CS model with the exchange interaction between $j = 3/2$ conduction electrons and a $J = 3/2$ local spin, the Kondo temperature $T^{(1)}_{\rm K}$ is defined by $j(j+1)/3\chi(T = 0)$. The $\chi(T)/\chi(0)$ ($\alpha = 1$) curve for the Γ_8 quartet is in agreement with that for the $J = 3/2$ spin completely, and we obtain $\chi(T^{(1)}_{\rm K})/\chi(0) \simeq 1/3$. As shown in Fig. 4, a crossover to the FL state appears with a $\ln T$ dependence in $\chi(T)$ for each value of α. We also obtain

$\chi(T_{\rm K}^{(0)})/\chi(0) \simeq 1/3$ for $\alpha = 0$, where $T_{\rm K}^{(0)}$ has been obtained from the logarithmic temperature dependent $\chi_{\rm Q}$ for $\alpha = 0$. Therefore we can define the Kondo temperature for arbitrary values of α using the equation $\chi(T_{\rm K}^{(\alpha)})/\chi(0) = 1/3$. Based on this analysis, we find that $T_{\rm K}^{(\alpha)}$ is analogous to $T_{\rm K}^{(1)}$ of the CS limit when $|\alpha|$ is close to unity. In Fig. 5, $\chi(0)$ is fitted well to the function $(T_{\rm K}^{(0)} + b\alpha^2)^{-1}$ from $\alpha = 1$ down to $\alpha \simeq 0.2$ for $\tilde{I}_{\rm o} = 0.2$ ($\alpha \simeq 0.4$ for $\tilde{I}_{\rm o} = 0.3$), where the constant b depends on $I_{\rm o}$. On the other hand, we find that $T_{\rm K}^{(\alpha)}$ is proportional to the inverse of this function in the same parameter region of α, and finally obtain

$$T_{\rm K}^{(\alpha)} \simeq 1.3(T_{\rm K}^{(0)} + cT_{\rm s}) \simeq \frac{1.3}{\chi(0)}, \tag{27}$$

where $c = 1.4$ for $\tilde{I}_{\rm o} = 0.2$ and $c = 0.86$ for $\tilde{I}_{\rm o} = 0.3$. This relation implies that the CS model can describe χ very well even if the $SU(4)$ symmetry is broken in real systems, since $|\alpha|$ is not usually so small. The strength of the anisotropy ($|\alpha| < 1$) is related to the reduction of the Kondo temperature. The large decrease of the quadrupolar coupling is seen in χ as a small difference from the CS limit. On the other hand, the crossover depending on the quadrupolar coupling can be observed more easily in $\chi_{\rm Q}$, since the observable crossover temperature $T_{\rm s}$ is very sensitive to the quadrupolar coupling.

IV. DISCUSSIONS

We have discussed the multipolar Kondo effect due to the Γ_8 quartet, based on the realistic Kondo model with the spherical tensor form. The exchange couplings depend on the crystal-field energy levels of the f^0, f^1 Γ_8 quartet, and f^2 Γ_1 singlet states. We can tune up the crystal field to leave the magnetic (dipolar + octupolar) exchange coupling $I_{\rm o}$ alone and obtain the new NFL state at low temperatures in this particle-hole symmetric case. This symmetry is usually broken by the nonmagnetic (quadrupolar + potential) coupling $I_{\rm e}$ in real systems. The latter coupling $I_{\rm e}$ behaves like a spin field in the 2CK model. The crossover from the NFL state to the FL state is related to the ratio of the two couplings α and we obtain two characteristic energy scales: one is the Kondo temperature $T_{\rm K}^{(\alpha)}$ reduced by the decrease of α from the CS limit ($\alpha = 1$), and the other is the small crossover temperature $T_{\rm s}$ ($\propto \alpha^2$) due to the NFL fixed point. The latter energy scale $T_{\rm s}$ clearly appears with the suppression of the logarithmic behavior in the quadrupolar susceptibility $\chi_{\rm Q}(T)$.

In our Kondo model, we have neglected the effect of the local excited states (Γ_3, Γ_4, Γ_5) in f^2, assuming that their energies are very large. If they are included, the Kondo model cannot be expressed with the spherical tensor form. The exchange process via the excited states makes the exchange interaction much more complicated, and prevents the total angular momentum from being a good quantum number if Γ_8 is mapped to $J = 3/2$. In addition, the twofold Γ_7 conduction electrons contribute to the exchange with the Γ_8 quartet and leads to a more complicated Kondo interaction. The larger number of degrees of freedom of the scattering process may stabilize the FL state more. For $\mathrm{Ce}_x\mathrm{La}_{1-x}\mathrm{B}_6$, we can expect a large crystal-field splitting between the ground and excited states in f^2 as well as between the Γ_8 and Γ_7 states in f^1. Since the f-electron is localized very well (the f-occupancy is close to one), the hybridization of the conduction band with the f-orbitals may be so small that the effect of the excited states becomes irrelevant. However, it is not clear which is a ground state for f^2 among the possible candidates Γ_1, Γ_3, and Γ_5.

Finally we discuss the connection between our results and the experiments for $\mathrm{Ce}_x\mathrm{La}_{1-x}\mathrm{B}_6$. According to the magnetic susceptibility and the resistivity, the Kondo temperature $T_{\rm K}$ is not sensitive to the Ce concentration x.[25,26] In addition, the x dependence of the lattice parameter is very small.[25] In our theory, $T_{\rm K}$ depends on the two Kondo couplings $I_{\rm o}$ and $I_{\rm e}$. As we discussed in the introduction, $T_{\rm Q}$ decreases more rapidly than $T_{\rm N}$ with decreasing x from $x = 1$. One can usually expect the quadrupolar order to be much sensitive to the lattice disorder since the quadrupolar moment is directly coupled to the lattice degrees of freedom. However, doping also changes the crystal field. It is possible that the quadrupolar RKKY coupling is suppressed more strongly by the change of x than the magnetic one, due to the different crystal-field dependence of the quadrupolar and magnetic exchange couplings of the conduction electrons with the Γ_8 quartets. We can conclude that the ratio of the two couplings $|I_{\rm e}/I_{\rm o}|$ decreases as x decreases from $x = 1$. Based on our results, T_K decreases with $|I_{\rm e}/I_{\rm o}|$, while it increases if $I_{\rm o}$ becomes larger. This is possible in general and thus we can obtain $T_{\rm K}$ to be less sensitive to doping than $T_{\rm Q}$ even if the Kondo couplings are varied.

In order to confirm our idea for the x dependence of $T_{\rm Q}$, we propose that the experimentalists measure χ and $\chi_{\rm Q}$ together for arbitrary x including the dilute limit ($x \ll 1$). $\chi_{\rm Q}$ is related to the elastic constant $(C_{11} - C_{12})/2$. It is very useful to observe the correlation between the two susceptibilities for the purpose of finding the local effect. When $|I_{\rm e}/I_{\rm o}| \simeq 1$, the correlation is expected to be linear in a temperature region where the Kondo effect is observed, since both χ and $\chi_{\rm Q}$ show almost the same temperature dependence. This linear feature tends to be lost as $|I_{\rm e}/I_{\rm o}|$ decreases form unity. Large deviations from the linear correlation can be found at low temperatures below $T_{\rm K}$. At higher temperatures, slight deviations are observable when $|I_{\rm e}|$ is very small. The same argument can be applied to the C_{44} mode as well. The amplitude of the quadrupolar Kondo coupling is also related to the type of quasiparticle excitations, particle-like or hole-like. If we are able to extract the single impurity effect, the thermopower measurement can detect the degree of particle-hole asymmetry which becomes smaller as the quadrupolar coupling decreases. It is also very interesting to search for the hidden NFL state related to the particle-hole symmetry not only in $\mathrm{Ce}_x\mathrm{La}_{1-x}\mathrm{B}_6$ but also in other cubic crystals including rare earth ions where a Γ_8 quartet is a ground state.

ACKNOWLEDGMENTS

The author thanks D. L. Cox and R. Shiina for stimulating discussions and critical reading of this manuscript. This research has been supported by JSPS Postdoctoral Fellowships for Research Abroad and by the US Department of Energy, Office of Science, Division of Materials Research.

---

[1] For a review see D. L. Cox and A. Zawadowski, Adv. Phys. **47**, 599 (1998) and references therein.

[2] A. C. Hewson, *The Kondo Problem to Heavy Fermions* (Cambridge University Press, 1993).

[3] K. Yamada, K. Hanzawa, and K. Yosida, Prog. Theor. Phys. Suppl. **108**, 141 (1992).

[4] H. Amitsuka and T. Sakakibara, J. Phys Soc. Jpn **63**, 736 (1994).

[5] S. Nishigori, K. Fujiwara, and T. Ito, Physica (Amsterdam) **B259-261**, 397 (1999).

[6] D. L. Cox, Physica (Amsterdam) **B186-188**, 312 (1993).

[7] M. C. Aronson, M. B. Maple, P. de Sa, A. M. Tsvelik, and R. Osborn, Europhys. Lett. **40**, 245 (1997).

[8] M. C. de Andrade, R. Chau, R. P. Dickey, N. R. Dilley, E. J. Freeman, D. A. Gajewski, M. B. Maple, R. Movshovich, A. H. Castro Neto, G. Castilla, and B. A. Jones, Phys. Rev. Lett. **81**, 5620 (1998).

[9] M. Koga, G. Zaránd, and D. L. Cox, submitted to Phys. Rev. Lett. (cond-mat/9903311).

[10] N. E. Bickers, D. L. Cox, and J. W. Wilkins, Physical Review B **36**, 2036 (1987).

[11] L. L. Hirst, Adv. Phys. **27**, 231 (1978).

[12] E. Pavarini and L. C. Andreani, Phys. Rev. B **56**, 5073 (1997).

[13] H. Kusunose and Y. Kuramoto, to be published in J. Phys. Soc. Jpn (cond-mat/9904292).

[14] M. Hiroi, S. Kobayashi, M. Sera, N. Kobayashi, and S. Kunii, J. Phys. Soc. Jpn **66**, 1762 (1997).

[15] T. Tayama, T. Sakakibara, K. Tenya, H. Amitsuka, and S. Kunii, J. Phys. Soc. Jpn **66**, 2268 (1997).

[16] J. R. Schrieffer and P. A. Wolff, Phys. Rev. **149**, 491 (1966).

[17] K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975); O. sakai, Y. Shimizu, and T. Kasuya, Prog. Theor. Phys. Suppl. **108**, 73 (1992).

[18] I. Affleck, A. W. W. Ludwig, H.-B. Pang, and D. L. Cox, Phys. Rev. B **45**, 7918 (1992).

[19] K. K. Lea, M. J. M. Leask, and W. P. Wolf, J. Phys. Chem. Solids **23**, 1381 (1962).

[20] Y. Onodera and M. Okazaki, J. Phys. Soc. Jpn **21**, 2400 (1966).

[21] T.-S. Kim, L. N. Oliveira, and D. L. Cox, Phys. Rev. B **55**, 12460 (1997).

[22] I Affleck, Nucl. Phys. B **336**, 517 (1990); I. Affleck and A. W. W. Ludwig, *ibid.* **352**, 849 (1991).

[23] M. Fabrizio, A. O. Gogolin, and P. Nozières, Phys. Rev. Lett. **74**, 4503 (1995).

[24] N. Andrei and A. Jerez, Phys. Rev. Lett. **74**, 4507 (1995).

[25] N. Sato, S. Kunii, I. Oguro, T. Komatsubara, and T. Kasuya, J. Phys. Soc. Jpn **53**, 3967 (1984).

[26] N. Sato, A. Sumiyama, S. Kunii, H. Nagano, and T. Kasuya, J. Phys. Soc. Jpn **54**, 1923 (1985).

(a)	q	j	(b)	Q	S	S_c	$El/\pi v_\mathrm{F}$
	1/2	0		0	1/2	0	0
	0	3/2		±1	0	1/2	1/8
	1/2	2		±2	1/2	0	1/2
				0	1/2	1	1/2
	1	3/2		±1	1	1/2	5/8
	3/2	0		0	3/2	0	1
	1/2	3		±2	1/2	1	1
				0	1/2	0	1
	1/2	1		0	1/2	1	1
	1/2	0		0	1/2	0	1

TABLE I. Lowest energy spectra at non-Fermi liquid fixed points: (a) For our model ($\alpha = 0$), q is an axial charge and j is a total angular momentum. (b) For the two-channel Kondo model, Q is a total number of particles measured from that of the ground state, S is a total spin, and S_c is an isospin associated with the two channels. The last column gives energies corresponding to the eigenstates for both (a) and (b), where we take the energy of the ground state as the origin.



FIG. 1. Crossover temperature. Over a wide range of the coupling ratio α, $T_{\rm s}^*$ is proportional to α^2.



FIG. 2. Temperature dependent quadrupolar susceptibility. From top to bottom, α is equal to 0, 0.04, 0.1, 0.2, 0.4, and 1.



FIG. 3. $T = 0$ values of quadrupolar susceptibility. For the small values of the coupling ratio α (< 0.2), $\chi_{\rm Q}$ is logarithmic in α.



FIG. 4. Temperature dependent magnetic susceptibility divided by each $T = 0$ value. From left to right, α is equal to 0, 0.2, 0.4, 0.6, 0.8, and 1.



FIG. 5. $T = 0$ values of magnetic susceptibility. When the coupling ratio α is not very small, χ is fitted to $(A + B\alpha^2)^{-1}$ (solid line), where $(A, B) = (2.7 \times 10^{-3}, 0.011)$ for $\tilde{I}_{\rm o} = 0.2$ and $(3.3 \times 10^{-3}, 0.049)$ for $\tilde{I}_{\rm o} = 0.3$.